advanced classical field theory

This page intentionally left blank

advanced classical field theory

giovanni giachetta
university of camerino, italy

luigi mangiarotti
university of camerino, italy

gennadi sardanashvily
moscow state university, russia

Published by

World Scientific Publishing Co. Pte. Ltd.

5 Toh Tuck Link, Singapore 596224

USA office: 27 Warren Street, Suite 401-402, Hackensack, NJ 07601

UK office: 57 Shelton Street, Covent Garden, London WC2H 9HE

British Library Cataloguing-in-Publication Data

A catalogue record for this book is available from the British Library.

ADVANCED CLASSICAL FIELD THEORY

Copyright © 2009 by World Scientific Publishing Co. Pte. Ltd.

All rights reserved. This book, or parts thereof, may not be reproduced in any form or by any means, electronic or mechanical, including photocopying, recording or any information storage and retrieval system now known or to be invented, without written permission from the Publisher.

For photocopying of material in this volume, please pay a copying fee through the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, USA. In this case permission to photocopy is not required from the publisher.

ISBN-13 978-981-283-895-7

ISBN-10 981-283-895-3

Printed in Singapore.

Preface

Contemporary quantum field theory is mainly developed as quantization of classical fields. Classical field theory thus is a necessary step towards quantum field theory. This book provides an exhaust mathematical foundation of Lagrangian classical field theory and its BRST extension for the purpose of quantization.

Lagrangian theory of Grassmann-graded (even and odd) fields on fibre bundles and graded manifolds is presented in the book in a very general setting. It is adequately formulated in geometric and algebraic topological terms of the jet manifolds and the variational bicomplex. The main ingredients in this formulation are cohomology of the variational bicomplex, the global first variational formula, variational symmetries and supersymmetries, the first Noether theorem, Noether identities, the direct and inverse second Noether theorems, and gauge symmetries.

Degenerate Lagrangian field theories are comprehensively investigated. The hierarchies of their non-trivial reducible Noether identities and gauge symmetries are described in homology terms. The relevant direct and inverse second Noether theorems are formulated in a very general setting.

The study of degeneracy of Lagrangian field theory straightforwardly leads to its BRST extension by Grassmann-graded antifields and ghosts which constitute the chain and cochain complexes of non-trivial Noether identities and gauge symmetries. In particular, a gauge operator is prolonged to a nilpotent BRST operator, and an original field Lagrangian is extended to a non-trivial solution of the classical master equation of Lagrangian BRST theory. This is a preliminary step towards quantization of classical Lagrangian field theory in terms of functional integrals.

The basic field theories, including gauge theory on principal bundles, gravitation theory on natural bundles, theory of spinor fields and topolog-

ical field theory, are presented in the book in a very complete way.

Our book addresses to a wide audience of theoreticians and mathematical physicists, and aims to be a guide to advanced differential geometric and algebraic topological methods in field theory.

With respect to mathematical prerequisites, the reader is expected to be familiar with the basics of differential geometry of fibre bundles. We have tried to give the necessary mathematical background, thus making the exposition self-contained. For the sake of convenience of the reader, several relevant mathematical topics are compiled in Appendixes.

Contents

<i>Preface</i>	v
<i>Introduction</i>	1
1. Differential calculus on fibre bundles	5
1.1 Geometry of fibre bundles	5
1.1.1 Manifold morphisms	6
1.1.2 Fibred manifolds and fibre bundles	7
1.1.3 Vector and affine bundles	12
1.1.4 Vector fields, distributions and foliations	18
1.1.5 Exterior and tangent-valued forms	21
1.2 Jet manifolds	26
1.3 Connections on fibre bundles	29
1.3.1 Connections as tangent-valued forms	30
1.3.2 Connections as jet bundle sections	32
1.3.3 Curvature and torsion	34
1.3.4 Linear connections	36
1.3.5 Affine connections	38
1.3.6 Flat connections	39
1.3.7 Second order connections	41
1.4 Composite bundles	42
1.5 Higher order jet manifolds	46
1.6 Differential operators and equations	51
1.7 Infinite order jet formalism	54
2. Lagrangian field theory on fibre bundles	61
2.1 Variational bicomplex	61

2.2	Lagrangian symmetries	66
2.3	Gauge symmetries	70
2.4	First order Lagrangian field theory	73
2.4.1	Cartan and Hamilton–De Donder equations . . .	75
2.4.2	Lagrangian conservation laws	78
2.4.3	Gauge conservation laws. Superpotential	80
2.4.4	Non-regular quadratic Lagrangians	83
2.4.5	Reduced second order Lagrangians	87
2.4.6	Background fields	88
2.4.7	Variation Euler–Lagrange equation. Jacobi fields .	90
2.5	Appendix. Cohomology of the variational bicomplex . . .	92
3.	Grassmann-graded Lagrangian field theory	99
3.1	Grassmann-graded algebraic calculus	99
3.2	Grassmann-graded differential calculus	104
3.3	Geometry of graded manifolds	107
3.4	Grassmann-graded variational bicomplex	115
3.5	Lagrangian theory of even and odd fields	120
3.6	Appendix. Cohomology of the Grassmann-graded varia- tional bicomplex	125
4.	Lagrangian BRST theory	129
4.1	Noether identities. The Koszul–Tate complex	130
4.2	Second Noether theorems in a general setting	140
4.3	BRST operator	147
4.4	BRST extended Lagrangian field theory	150
4.5	Appendix. Noether identities of differential operators . . .	154
5.	Gauge theory on principal bundles	165
5.1	Geometry of Lie groups	165
5.2	Bundles with structure groups	169
5.3	Principal bundles	171
5.4	Principal connections. Gauge fields	175
5.5	Canonical principal connection	179
5.6	Gauge transformations	181
5.7	Geometry of associated bundles. Matter fields	184
5.8	Yang–Mills gauge theory	188
5.8.1	Gauge field Lagrangian	188

5.8.2	Conservation laws	190
5.8.3	BRST extension	192
5.8.4	Matter field Lagrangian	194
5.9	Yang–Mills supergauge theory	196
5.10	Reduced structure. Higgs fields	198
5.10.1	Reduction of a structure group	198
5.10.2	Reduced subbundles	200
5.10.3	Reducible principal connections	202
5.10.4	Associated bundles. Matter and Higgs fields . . .	203
5.10.5	Matter field Lagrangian	207
5.11	Appendix. Non-linear realization of Lie algebras	211
6.	Gravitation theory on natural bundles	215
6.1	Natural bundles	215
6.2	Linear world connections	219
6.3	Lorentz reduced structure. Gravitational field	223
6.4	Space-time structure	228
6.5	Gauge gravitation theory	232
6.6	Energy-momentum conservation law	236
6.7	Appendix. Affine world connections	238
7.	Spinor fields	243
7.1	Clifford algebras and Dirac spinors	243
7.2	Dirac spinor structure	246
7.3	Universal spinor structure	252
7.4	Dirac fermion fields	258
8.	Topological field theories	263
8.1	Topological characteristics of principal connections	263
8.1.1	Characteristic classes of principal connections . .	264
8.1.2	Flat principal connections	266
8.1.3	Chern classes of unitary principal connections . .	270
8.1.4	Characteristic classes of world connections	274
8.2	Chern–Simons topological field theory	278
8.3	Topological BF theory	283
8.4	Lagrangian theory of submanifolds	286
9.	Covariant Hamiltonian field theory	293

9.1	Polysymplectic Hamiltonian formalism	293
9.2	Associated Hamiltonian and Lagrangian systems	298
9.3	Hamiltonian conservation laws	304
9.4	Quadratic Lagrangian and Hamiltonian systems	306
9.5	Example. Yang–Mills gauge theory	313
9.6	Variation Hamilton equations. Jacobi fields	316
10.	Appendixes	319
10.1	Commutative algebra	319
10.2	Differential operators on modules	324
10.3	Homology and cohomology of complexes	327
10.4	Cohomology of groups	330
10.5	Cohomology of Lie algebras	333
10.6	Differential calculus over a commutative ring	334
10.7	Sheaf cohomology	337
10.8	Local-ring spaces	346
10.9	Cohomology of smooth manifolds	348
10.10	Leafwise and fibrewise cohomology	354
	<i>Bibliography</i>	359
	<i>Index</i>	369

Introduction

Contemporary quantum field theory is mainly developed as quantization of classical fields. In particular, a generating functional of Green functions in perturbative quantum field theory depends on an action functional of classical fields. In contrast with quantum field theory, classical field theory can be formulated in a strict mathematical way.

Observable classical fields are an electromagnetic field, Dirac spinor fields and a gravitational field on a world real smooth manifold. Their dynamic equations are Euler–Lagrange equations derived from a Lagrangian. One also considers classical non-Abelian gauge fields and Higgs fields. Basing on these models, we study Lagrangian theory of classical Grassmann-graded (even and odd) fields on an arbitrary smooth manifold in a very general setting. Geometry of principal bundles is known to provide the adequate mathematical formulation of classical gauge theory. Generalizing this formulation, we define even classical fields as sections of smooth fibre bundles and, accordingly, develop their Lagrangian theory as Lagrangian theory on fibre bundles.

Note that, treating classical field theory, we are in the category of finite-dimensional smooth real manifolds, which are Hausdorff, second-countable and paracompact. Let X be such a manifold. If classical fields form a projective $C^\infty(X)$ -module of finite rank, their representation by sections of a fibre bundle follows from the well-known Serre–Swan theorem.

Lagrangian theory on fibre bundles is adequately formulated in algebraic terms of the variational bicomplex of exterior forms on jet manifolds [3; 17; 59]. This formulation is straightforwardly extended to Lagrangian theory of even and odd fields by means of the Grassmann-graded variational bicomplex [9; 14; 59]. Cohomology of this bicomplex provides the global first variational formula for Lagrangians and Euler–Lagrange operators, the first

Noether theorem and conservation laws in a general case of supersymmetries depending on derivatives of fields of any order.

Note that there are different descriptions of odd fields on graded manifolds [27; 118] and supermanifolds [29; 45]. Both graded manifolds and supermanifolds are described in terms of sheaves of graded commutative algebras [10]. However, graded manifolds are characterized by sheaves on smooth manifolds, while supermanifolds are constructed by gluing of sheaves on supervector spaces. Treating odd fields on a smooth manifold X , we follow the Serre–Swan theorem generalized to graded manifolds [14]. It states that, if a Grassmann $C^\infty(X)$ -algebra is an exterior algebra of some projective $C^\infty(X)$ -module of finite rank, it is isomorphic to the algebra of graded functions on a graded manifold whose body is X .

Quantization of Lagrangian field theory essentially depends on its degeneracy characterized by a family of non-trivial reducible Noether identities [9; 15; 63]. A problem is that any Euler–Lagrange operator satisfies Noether identities which therefore must be separated into the trivial and non-trivial ones. These Noether identities can obey first-stage Noether identities, which in turn are subject to the second-stage ones, and so on. If certain conditions hold, this hierarchy of Noether identities is described by the exact Koszul–Tate chain complex of antifields possessing the boundary operator whose nilpotentness is equivalent to all non-trivial Noether and higher-stage Noether identities [14; 15].

The inverse second Noether theorem formulated in homology terms associates to this Koszul–Tate complex the cochain sequence of ghosts with the ascent operator, called the gauge operator, whose components are non-trivial gauge and higher-stage gauge symmetries of Lagrangian field theory [15]. These gauge symmetries are parameterized by odd and even ghosts so that k -stage gauge symmetries act on $(k - 1)$ -stage ghosts.

It should be emphasized that the gauge operator unlike the Koszul–Tate one is not nilpotent, unless gauge symmetries are Abelian. Gauge symmetries are said to be algebraically closed if this gauge operator admits a nilpotent extension where k -stage gauge symmetries are extended to k -stage BRST (Becchi–Rouet–Stora–Tyutin) transformations acting both on $(k - 1)$ -stage and k -stage ghosts [61]. This nilpotent extension is called the BRST operator. If the BRST operator exists, the cochain sequence of ghosts is brought into the BRST complex.

The Koszul–Tate and BRST complexes provide a BRST extension of original Lagrangian field theory. This extension exemplifies so called field-antifield theory whose Lagrangians are required to satisfy a certain con-

dition, called the classical master equation. An original Lagrangian is extended to a proper solution of the master equation if the BRST operator exists [15]. This extended Lagrangian, dependent on original fields, ghosts and antifields, is a first step towards quantization of classical field theory in terms of functional integrals [9; 63].

The basic field theories, including gauge theory on principal bundles (Chapter 5), gravitation theory on natural bundles (Chapter 6), theory of spinor fields (Chapter 7) and topological field theory (Chapter 8) are presented in the book in a complete way.

The reader also can find a number of original topics, including: general theory of connections (Section 1.3), geometry of composite bundles (Section 1.4), infinite-order jet formalism (Section 1.7), generalized symmetries (Section 2.2), Grassmann-graded Lagrangian field theory (Section 3.5), second Noether theorems in a general setting (Section 4.2), the BRST complex (Section 4.3), classical Higgs field theory (Section 5.10), gauge theory of gravity as a Higgs field (Section 6.5), gauge energy-momentum conservation laws (Section 6.6), composite spinor bundles (Section 7.3), global Chern–Simons topological field theory (Section 8.2), topological BF (background field) theory (Section 8.3), covariant Hamiltonian field theory (Chapter 9).

For the sake of convenience of the reader, several relevant mathematical topics are compiled in Chapter 10.