Serious limitations of the strong equivalence principle

De-Chang Dai

*Institute of Natural Sciences, Shanghai Key Laboratory for Particle Physics and Cosmology,
and Center for Astrophysics and Astronomy, Department of Physics and Astronomy,
Shanghai Jiao Tong University, Shanghai 200240, China
diedachung@gmail.com*

Received 4 March 2017

Accepted 30 March 2017

Published 7 April 2017

It is well known that an accelerated charged particle radiates away energy. However, whether an accelerated neutral composite particle radiates away energy is unclear. We study decoherent Larmor radiation from an accelerated neutral composite object. We find that the neutral object's long wavelength radiation is highly suppressed because radiation from different charges is canceled out. However, the neutral object radiates high energy or short wavelength radiation without any suppression. In that case, radiation from each particle can be treated independently, and it is called the decoherent radiation. We compare a hydrogen atom's decoherent Larmor radiation with its gravitational radiation while the atom is in a circular orbit around a star. Gravitational radiation is stronger than the electromagnetic radiation if the orbital radius is larger than some critical radius. Since the decoherent radiation is related to the object's structure, this implies that the strong equivalence principle which states that gravitational motion does not depend on an object's constitution has severe limitations.

Keywords: Equivalence principle; coherent Larmor radiation; acceleration.

PACS numbers: 04.20.Cv, 04.30.Db, 04.62.+v, 03.50.De

1. Introduction

Since Galileo discovered that objects of any weight and composition fall toward the Earth at the same rate, the equivalence principle has played an important role in most of the theories of gravity. Einstein formulated the General Relativity based on the acceleration equivalence principle. As a consequence, a phenomenon observed in an accelerated system might also be observed in a gravitational field, e.g. Hawking radiation and Unruh radiation.^{1,2}

This is an Open Access article published by World Scientific Publishing Company. It is distributed under the terms of the Creative Commons Attribution 4.0 (CC-BY) License. Further distribution of this work is permitted, provided the original work is properly cited.

There is no doubt that gravity and acceleration share common physical features. But, Galileo's original observation that objects of any weight and composition respond to gravity equally is still subject to questioning. It is known that an accelerated charged particle radiates away energy through the Larmor radiation. This implies that a charged particle will emit radiation under gravitational acceleration and its trajectory is not going to be the same as a neutral particle's. The charged particle's equation of motion satisfies Abraham–Lorentz–Dirac equation in the curved space–time instead of the simple geodesic path.^{3–6}

Since the free falling objects' trajectories depend on their charges, Galileo's original idea must be modified. However, most of the falling objects are not charged. So far only the accelerated charged particles were studied in this context.^{7,8} However, if the composite particle is neutral, the individual charges cancel out. It is still not clear whether a neutral composite object follows a trajectory different from the geodesic path. It is also unknown whether a neutral composite object radiates away energy, since the radiation might allow an observer to reconstruct the distribution of matter which made a black hole.^{9,10}

To study this question, we first note that a charged particle is different from a neutral particle, because it radiates away electromagnetic fields. Therefore, if one can study radiation, then he could reconstruct the distribution of matter in a composite object. Similar problem has been studied in the synchrotron radiation theoretically and experimentally.^{11–16} In general, electrons are accelerated in groups in a synchrotron. It is found that each electron radiates away electromagnetic field as a single charged particle if the radiation wavelength is shorter than the distance between electrons in the electron bundle. This is called decoherent radiation. Charged particles in that case are independent from each other and must be treated independently. However, if the electromagnetic field's wavelength is longer than the size of the electron bundle, the radiation is the same as a single charged particle with charge Ne (N is the number of electrons). The radiation is highly enhanced and is called the coherent radiation.

Based on the coherent radiation theory, radiation from a neutral composite object is highly suppressed for long wavelength mode because the object must be treated as a coherent single object. But, the object can radiate away short wavelength modes because these modes are radiated by all the components independently. We study this coherent radiation and decoherent radiation with the use of an interference function $f(\omega)$. If $f(\omega) \rightarrow 0$, the radiation is coherent. We find that whether the radiation is coherent or decoherent depends on the object's size. For a charged particle pair with distance l , the critical energy is $\omega_c \approx 1/l$. For a hydrogen atom (its radius is about a_0), the critical energy is $\omega_c \approx 1/a_0$. We compare an accelerated hydrogen atom's electromagnetic radiation to its gravitational radiation, while the atom is in a circular orbit around a star with mass equal to $M_\odot$, and find that the gravitational radiation is stronger than the electromagnetic radiation if its orbital radius is larger than some critical radius. In the following, we

introduce the coherent radiation theory and use a charged particle pair as a simple example. Then, we calculate the hydrogen atom's interference function.

2. Coherent Radiation from a Multiparticle System

It is known that an accelerated charged particle radiates away energy, which is called Larmor radiation. Its intensity distribution is¹⁷

$$\frac{d^2 I}{d\omega d\Omega} = \frac{\omega^2 e^2}{16\pi^3} \left| \int \hat{n} \times (\hat{n} \times \boldsymbol{\beta}) e^{i\omega(t-\hat{n}\cdot\mathbf{r})} dt \right|^2. \quad (1)$$

Here, $\boldsymbol{\beta}$ is the velocity of light ($\mathbf{v}/c$), $\hat{n}$ is the direction from the charged particle to an observer at infinity, $\mathbf{r}$ is the location of the particle. ω is the energy of the particle, e is the particle's charge and Ω is the solid angle. We set $c = \hbar = 1$. In general, an object is made of several particles which have their own charges. The radiation comes from each charged particle

$$\frac{d^2 I}{d\omega d\Omega} = \frac{\omega^2 e^2}{16\pi^3} \left| \int \hat{n} \times (\hat{n} \times \mathbf{J}) e^{i\omega(t-\hat{n}\cdot\mathbf{r})} dt dx^3 \right|^2. \quad (2)$$

$\mathbf{J} = \rho\boldsymbol{\beta}$, ρ is the charge density. Each particle's radiation interferes with radiation from other particles. The object's center-of-mass is at $\mathbf{r}_c$. $\mathbf{r}$ can be transferred to the center-of-mass frame, $\mathbf{r}'$, with the relation $\mathbf{r} = \mathbf{r}_c - \mathbf{r}'$. Then, the radiation can be written as¹²

$$\frac{d^2 I}{d\omega d\Omega} = \frac{\omega^2 e^2}{16\pi^3} f(\omega) \left| \int \hat{n} \times (\hat{n} \times \boldsymbol{\beta}) e^{i\omega(t-\hat{n}\cdot\mathbf{r}_c)} dt \right|^2, \quad (3)$$

$$f(\omega) = \left| \int \rho e^{i\omega\hat{n}\cdot\mathbf{r}'} dx'^3 \right|^2. \quad (4)$$

Equation (3) is similar to a single particle Larmor radiation multiplied by an interference factor $f(\omega)$. Therefore, if $f(\omega)$ is obtained, then the radiation is known. In the following, we will focus on calculating $f(\omega)$.

3. A Charged Particle Pair

The simplest composite object is made of a charged particle pair. We assume that there is one particle with charge q located at $(0, 0, l/2)$ and another particle with charge $-q$ located at $(0, 0, -l/2)$ in the center-of-mass frame (as in Fig. 1). The center-of-mass is located at $\mathbf{r}_c$. While in general particles may be free to move with respect to each other, we calculate only instantaneous radiation and keep the structure fixed. The charge density can be written with two delta functions,

$$\rho = \delta\left(\mathbf{r} - \frac{l}{2}\hat{z}\right) - \delta\left(\mathbf{r} + \frac{l}{2}\hat{z}\right). \quad (5)$$

Fig. 1. There are two charged particles in the system. One particle with charge q is at $(0, 0, l/2)$ and the other particle with charge $-q$ is at $(0, 0, -l/2)$.

To simplify the study, we consider the nonrelativistic case, and the Lorentz contraction is neglected. The interference function $f(\omega)$ is

$$f(\omega) = \left| e^{i\omega \frac{l}{2} \hat{n} \cdot \hat{z}} - e^{-i\omega \frac{l}{2} \hat{n} \cdot \hat{z}} \right|^2 = 4 \sin^2 \left(\omega \frac{l}{2} \hat{n} \cdot \hat{z} \right). \quad (6)$$

This is the same as double slit interference with phase difference π . If the observer is at $(0, 0, \infty)$, modes with $\omega \ll 1/l$ cancel out each other and $f(\omega) \rightarrow 0$. The interference pattern appears for $\omega l \gg 1$ (Fig. 2). The intensity of the interference is between 0 and 4. The oscillation cycle is $\Delta\omega = \frac{4\pi}{l}$. While ω is large enough, a detector will not be able to pick up the oscillation pattern and it will record only the average effect. On average, the intensity is $\bar{f}(\omega) = 2$. This value is

Fig. 2. The interference function for two charged particles, $f(\omega)$, as a function of ω . For low ω , radiation from these two particles is coherent and cancel out each other. The observer is in the direction of the z -axis. For high ω modes, the interference pattern appears. It oscillates between 0 and 4 and the oscillation period is $\Delta\omega = \frac{4\pi}{l}$. On average, it becomes equal to 2. For a finite size detector, one will not be able to distinguish the oscillations and it is as if the radiation is coming from two independent charged particles.

Fig. 3. There are four different interference functions, $f(\omega)$, as a function of ωl in the figure. The x -axis is the angle between the observer's direction and z -axis, $\varphi = \cos^{-1}(\hat{n} \cdot \hat{z})$. For $\omega l \ll 1$, $f(\omega)$ is almost 0, because radiation from these two charged particles cancel out. For $\omega l \gg 1$, the interference pattern starts to appear and its period becomes shorter and shorter with respect to φ . When a detector cannot distinguish the oscillation, the radiation is identical to radiation from two independent particles.

what one expects from two independent charged particles. Therefore, even though the object's total charge is 0, the high energy radiation is the same as the radiation from two independent particles.

For an observer at different location, the interference can be described by the angle $\varphi = \cos^{-1}(\hat{n} \cdot \hat{z})$. Again, the interference function, $f(\omega)$, is canceled out for $\omega \ll 1/l$ and the interference pattern appears while $\omega \gtrsim 1/l$ (Fig. 3). The oscillation pattern is between 0 and 4 and oscillation changes much quicker for larger ω . Its average is $\bar{f}(\omega) = 2$. This proves that the radiation with shorter wavelength acts as if it is coming from two independent charged particles or in other words, two charged particles radiate away energy decoherently.

4. Hydrogen

A neutral atom has no net charge. At first glimpse, it seems that an accelerated atom will not radiate any electromagnetic wave. However, since an atom has its own structure, its high ω radiation should not be canceled out. This can be tested by calculating the interference function $f(\omega)$. To make it simple, we will focus on the simplest atom, hydrogen. A hydrogen has a proton in the center and an electron circulating around it. The acceleration will distort the hydrogen's wave function.¹⁸ Here, we focus on small acceleration and assume that the distortion can be neglected. We also consider low velocities so that the Lorentz distortion is

neglected. The electron's wave function in the center-of-mass frame is

$$\phi_{n,l,m}^0(r, \theta, \phi) = R_{n,l} Y_l^m(\theta, \phi), \quad (7)$$

$$R_{n,l} \sim \exp\left(-\frac{r}{na_0}\right) \left(\frac{r}{na_0}\right)^l L_{n-l-1}^{2l+1}\left(\frac{2r}{na_0}\right). \quad (8)$$

$L_{n-l-1}^{2l+1}(x)$ is the associated Laguerre polynomial. $Y_l^m(\theta, \phi)$ is the spherical harmonic function of degree l and order m . a_0 is the Bohr radius, $a_0 = \frac{1}{me^2} = 5.2917721067 \times 10^{-11}$ m. The ground and first excited states' wave functions are

$$\phi_{1,0,0}^0 = \left(\frac{1}{\pi a_0^3}\right)^{\frac{1}{2}} \exp\left(-\frac{r}{a_0}\right), \quad (9)$$

$$\phi_{2,0,0}^0 = \left(\frac{1}{32\pi a_0^3}\right)^{\frac{1}{2}} \left(2 - \frac{r}{a_0}\right) \exp\left(-\frac{r}{2a_0}\right), \quad (10)$$

$$\phi_{2,1,0}^0 = \left(\frac{1}{32\pi a_0^3}\right)^{\frac{1}{2}} \frac{r}{a_0} \exp\left(-\frac{r}{2a_0}\right) \cos \theta, \quad (11)$$

$$\phi_{2,1,\pm 1}^0 = \mp \left(\frac{1}{64\pi a_0^3}\right)^{\frac{1}{2}} \frac{r}{a_0} \exp\left(-\frac{r}{2a_0}\right) \sin \theta e^{\pm i\phi}. \quad (12)$$

The electron's probability function is obtained from its wave function

$$\rho_e = \phi^* \phi \quad (13)$$

and the proton can be treated as a point particle at the center

$$\rho_p = \delta(\mathbf{r}). \quad (14)$$

An electron has a negative charge and a proton has a positive charge, so the total charge distribution is

$$\rho = \rho_p - \rho_e. \quad (15)$$

The interference function can be calculated directly from Eq. (4). Figure 4 shows the interference function of s -wave states ($l = 0$, $m = 0$). $f(\omega)$ is canceled out for small ω . While ω increases, $f(\omega)$ increases and achieves 1 at very large ω . The higher n states become decoherent at lower ω than the lower n states did because the higher n states have larger radii. However, the decoherent radiation is not a combination of the proton and electron. The electron's radiation is canceled out by itself, because of its own density distribution. Only the proton's radiation is left. The atom radiates high ω modes as if it is a single proton.

If the acceleration does not change very quick, there will not be much high frequency radiation ($\tilde{\beta}(\omega) \rightarrow 0$ as $\omega a_0 \rightarrow \infty$). In that case, we find the ground state's $f(\omega)$ is

$$f(\omega) = \left| \int \rho e^{i\omega \hat{n} \cdot \mathbf{r}_e} d\mathbf{x}^3 \right|^2 = \frac{\omega^4 a_0^4 (\omega^2 a_0^2 + 8)^2}{(16 + 8\omega^2 a_0^2 + \omega^4 a_0^4)^2}. \quad (16)$$

Fig. 4. The four curves are the interference functions for different quantum states, $(n, 0, 0)$. Since the larger n state has a larger radius, radiation becomes decoherent for much smaller ω . Since all of the four quantum states are spherically symmetric, the observer's direction does not affect the interference pattern.

In a low ω approximation,

$$f(\omega) \approx \frac{\omega^4 a_0^4}{4}. \quad (17)$$

We consider the nonrelativistic approximation. The velocity is much lower than the speed of light, $\beta \ll 1$

$$\left| \int \hat{n} \times (\hat{n} \times \beta) e^{i\omega(t-\hat{n} \cdot \mathbf{r})} dt \right|^2 \approx \left| \int \sin(\alpha) \beta e^{i\omega t} dt \right|^2 = \sin^2(\alpha) \tilde{\beta}^2. \quad (18)$$

Here, $\tilde{\beta}$ is β 's Fourier transform,

$$\tilde{\beta} = \int \beta e^{i\omega t} dt. \quad (19)$$

From Eq. (3), the intensity is

$$\frac{d^2 I}{d\omega d\Omega} = \frac{\omega^2 e^2}{16\pi^3} \frac{\omega^4 a_0^4}{4} \sin^2(\alpha) \tilde{\beta}^2. \quad (20)$$

The total released energy is

$$I = \int \frac{\omega^2 e^2}{16\pi^3} \frac{\omega^4 a_0^4}{4} \sin^2(\alpha) \tilde{\beta}^2 d\omega d\Omega = \int_0^\infty \frac{e^2 \omega^6 a_0^4}{24\pi^2} \tilde{\beta}^2 d\omega = \int_{-\infty}^\infty \frac{e^2 a_0^4}{24\pi} (\partial_t^3 \beta)^2 dt. \quad (21)$$

This implies that the energy flux is $\frac{e^2 a_0^4}{24\pi} (\partial_t^3 \beta)^2$, if the acceleration is changing slowly.

5. Comparing Neutral Atom's Electromagnetic and Gravitational Radiation

It is known that an accelerated particle also radiates away gravitational waves. A freely falling composite object may radiate both gravitational waves and electromagnetic waves. To simplify the case, we study a composite particle under the gravitational influence of a heavy object with mass M . We consider a circular orbit. Then, the average power of gravitational radiation over one period of the motion is¹⁹

$$\left\langle \frac{dI_g}{dt} \right\rangle = \frac{32G^4}{5} \frac{M^2 m^2 (M + m)}{R^5}. \quad (22)$$

Here, R is the orbital radius and m is the composite particle's mass. In this study, we assume that the composite particle is a hydrogen atom in this ground state. The orbital angular frequency is $\omega_o^2 = \frac{GM}{R^3}$.

The composite particle's electromagnetic radiation can be found from Eq. (21),

$$\frac{dI}{dt} = \frac{e^2 a_0^4 R^2 \omega_o^8}{24\pi} = \frac{e^2 a_0^4 G^4 M^4}{24\pi R^{10}}. \quad (23)$$

One can compare Eq. (22) with Eq. (23). The gravitational radiation is stronger than the electromagnetic radiation if its orbital radius is larger than the critical radius,

$$R_c = \left(\frac{5e^2 a_0^4 M}{768\pi m^2} \right)^{1/5}. \quad (24)$$

If the heavy object is of the sun's mass ($M = 1M_\odot$), the critical radius is $R_c = 1.96$ m. This is much smaller than the sun's radius. Therefore, in the sun's case, a freely falling hydrogen atom still radiates away much more gravitational than electromagnetic energy. However, this may change for a very massive and compact object.

6. Conclusion

We studied the decoherent Larmor radiation from a neutral object which is made of charged particles. This radiation depends on the object's structure. For a charged particle pair separated by the distance l , the characteristic energy is $\omega_c \approx 1/l$. The object's low ω radiation ($\omega \ll \omega_c$) must be treated as if it is coming from a single object and radiation is highly suppressed. The object's high ω modes ($\omega \gg \omega_c$) must be treated as if they are coming from independent particles. They are radiated decoherently.

The same argument is applied to the hydrogen atom. A hydrogen atom's radius is about a_0 . Its characteristic energy is $\omega_c \approx 1/a_0$. The $\omega \ll \omega_c$ modes are highly suppressed because the charged particles radiate coherently. The $\omega \gg \omega_c$ modes are radiated decoherently and one can treat the electron and proton independently. However, at this energy scale, electrons cannot be treated as single independent

particles. Electron's radiation is canceled out because of electron's charge density distribution. The high ω modes are mainly from the proton. The radiation is similar to the radiation from a single charged particle, instead of the radiation from two charged particles. If the acceleration does not change very quickly, the high ω radiation is weak. In this case, the total radiation depends on the third time derivative of velocity (Eq. (21)).

An accelerated neutral atom emits not only electromagnetic waves, but also gravitational waves. Both of these processes are highly suppressed. The electromagnetic radiation is suppressed by charge cancellation while the gravitational radiation is suppressed by the weak gravitational coupling. We compare these two types of radiations for a hydrogen atom orbiting a star of solar mass. We find that the gravitational radiation is much stronger than the electromagnetic radiation if the orbital radius is larger than $R_c = 1.96$ m.

This result shows that an accelerated particle will radiate energy according to its structure. This radiation may be highly suppressed because of the constituent particles' interference, but it is not fully canceled out. Therefore, if we drop two objects with different internal structures in a gravitational field, these two objects will radiate away different amounts of energy and will fall at a different rate. The strong equivalence principle which states that gravitational motion does not depend on an object's constitution must be revised. It is also possible to reconstruct the falling matter's internal structure through the radiation.⁹

Acknowledgments

De-Chang Dai was supported by the National Science Foundation of China (grant Nos. 11433001 and 11447601), National Basic Research Program of China (973 Program 2015CB857001), No. 14ZR1423200 from the Office of Science and Technology in Shanghai Municipal Government, the key laboratory grant from the Office of Science and Technology in Shanghai Municipal Government (No. 11DZ2260700) and the Program of Shanghai Academic/Technology Research Leader under grant No. 16XD1401600.

References

1. W. G. Unruh, *Phys. Rev. D* **14**, 870 (1976), doi:10.1103/PhysRevD.14.870.
2. J. B. Hartle and S. W. Hawking, *Phys. Rev. D* **13**, 2188 (1976), doi:10.1103/PhysRevD.13.2188.
3. M. Abraham and R. Becker, *Electricity and Magnetism* (Blackie, London, 1937).
4. P. A. M. Dirac, *Proc. R. Soc. A* **167**, 148 (1938).
5. H. A. Lorentz, *The Theory of Electrons* (Dover Books, New York, 1952).
6. B. S. DeWitt and R. W. Brehme, *Ann. Phys.* **9**, 220 (1960).
7. F. Rohrlich, *Amer. J. Phys.* **34**, 1206 (1966).
8. D. G. Boulware, *Ann. Phys.* **124**, 169 (1980).
9. D. C. Dai and D. Stojkovic, *Phys. Lett. B* **758**, 412 (2016), arXiv:1605.06026.
10. D. C. Dai and D. Stojkovic, *Phys. Lett. B* **758**, 429 (2016), doi:10.1016/j.physletb.2016.05.037, arXiv:1601.07921 [gr-qc].

11. J. S. Nodvick and D. S. Saxon, *Phys. Rev.* **96**, 180 (1954).
12. C. J. Hirschmugl, M. Sagurton and G. P. Williams, *Phys. Rev. A* **44**, 1316 (1991).
13. G. L. Carr, M. C. Martin, W. R. McKinney, K. Jorda, G. R. Neil and G. P. Williams, *Nature* **420**, 153 (2002), doi:10.1038/nature01175.
14. G. L. Orlandi, *Opt. Commun.* **211**, 109 (2002).
15. G. L. Orlandi, *Opt. Commun.* **267**, 322 (2006).
16. M. Hosaka *et al.*, *Phys. Rev. Spec. Top. Accel. Beams* **16**, 020701 (2013).
17. J. D. Jackson, *Classical Electrodynamics*, 3rd edn. (John Wiley & Sons, New York, 1999).
18. D. C. Dai, *Phys. Lett. A* **380**, 3601 (2016), doi:10.1016/j.physleta.2016.08.056.
19. P. C. Peters and J. Mathews, *Phys. Rev.* **131**, 435 (1963).