ERRATUM

MAGNETIC FIELD AND CURRENT DEPENDENCE OF
AVERAGE ACTIVATION ENERGY IN YBCO HIGH- T_c
GRANULAR SUPERCONDUCTORS

[*Mod. Phys. Lett. B*, Vol. 15, No. 16 (2001) 555–559]

V. POPESCU* and A. STEPANESCU†

**Department of Physics, University “Politehnica” Bucharest,
Splaiul Independentei 313, 77206 Bucharest, Romania*

†*Dip. Fisica, Politecnico di Torino, C. so Duca Abruzzi 24, 10129 Torino, Italy*

In Sec. 2, Results and Discussion, the incorrect figures were given. The correct figures should have been:

Fig. 1. The temperature dependence of the noise spectral power density RN at 10 Hz and 75 mA for two magnetic fields ($H = 120$ Oe and $H = 757$ Oe). When the magnetic field is increased, the noise peak shifts towards lower temperatures. The field dependence of the activation energy U is well approximated by $U(H) \sim H^{-0.17}$.

Fig. 2. The temperature dependence of the noise spectral power density RN at 10 Hz and 436 Oe for two values of the bias current ($I = 75.3$ mA and $I = 100$ mA). When the bias current is increased, the noise peak shifts towards lower temperatures. The current dependence of the activation energy U is well approximated by $U(I) \sim I^{-3/2}$.